

Cervical cancer in England 1996 to 2004

Mortality trends

Unit of Health Care Epidemiology,
Oxford University, and
South East England Public Health Observatory
2006



Cervical cancer in England 1996 to 2004.

Mortality trends

**Authors: Michael Goldacre, Marie Duncan, Paula Cook-Mozaffari,
Matthew Davidson, Henry McGuinness, Daniel Meddings**

**Published by: Unit of Health-Care Epidemiology, Oxford University, and
South-East England Public Health Observatory, 2006**

This document provides a profile of trends in mortality for cervical cancer in England. The period covered is January 1 1996 to December 31 2004. The data are analysed from mortality files supplied to the South East England Public Health Observatories (SEPHO) by the Office for National Statistics (ONS). Mortality rates were calculated for the condition certified as the underlying cause of death and for the disease certified as any mention on the death certificates. Age-specific mortality rates in 5-year age-groups were used to calculate age-standardised mortality rates with 95% confidence intervals for each time period. Age-truncated rates, with age-standardisation within them, are presented. We calculated the average annual percentage change from 1996-2004 in mortality rates for mentions, using data for each individual calendar year, by fitting linear regression models to the logarithms of the death rates. Age-standardised rates for each individual calendar year were calculated by applying the age-specific rates in each five-year age-group in each calendar year to the European standard population.

The Table shows age- and sex-specific mortality rates per million resident population. The Graph shows age-standardised mortality rates for mortality rates based on underlying cause and for mortality rates based on mentions.

Some technical notes:

- Dataset used: UHCE-built multi-year file for England, England_sepho.deaths.reg96-04unf300.v2, using the SEPHO mortality files for individual calendar years supplied by the ONS
- Deaths were coded to ICD9 180 for the years 1996-2000 and to ICD10 C53 for 2001-2004
- Diagnosis position: underlying cause, and 'all mentions'
- Period covered: deaths by date of occurrence from 1.1.1996 to 31.12.2004. Deaths in the final year for which data are available, namely 2004, may be slightly low due to late registration of some deaths
- Age/sex restriction: all ages, males and females separately
- Restrictions in record selection: none
- Place of residence at time of death: England

We are very grateful for computing support from Myfanwy Griffith and Leicester Gill. Work funded by the NHS National Centre for Research Capacity Development. Henry McGuinness and Daniel Meddings are funded by the South East England Public Health Observatory. Ethical approval: REC references 04/Q2006/175 and 04/Q2006/176.

Copyright © 2006 UHCE, Oxford University

Cervical cancer in England

Number of deaths, age-standardised mortality rates 1996-2004 for underlying cause and for mentions; average annual change in rates for mentions, and 95% confidence intervals on the change in mentions

Age group	<i>n</i>	Rates per million		Average annual % change	(95% CI)	
		Underlying cause 1996-04	Mentions 1996-04			
Women						
0-34	515	4.9	4.9	-3.3	-7.1	0.7
35-44	1202	40.6	42.3	-9.3	-11.3	-7.4
45-54	1512	55.1	58.7	-3.9	-5.3	-2.6
55-64	1567	67.5	74.9	-4.6	-6.5	-2.7
65-74	1967	93.5	109.2	-5.3	-6.5	-4.2
75+	3619	160.5	191.6	-3.4	-4.3	-2.5
all age	10382	36.2	40.2	-5.0	-5.7	-4.4

Cervical cancer in England: age-standardised mortality rates 1996-2004, females

